

**EC FERTILIZER**  
**Potassium Sulfate containing Magnesium Salt**  
 Performans PotasMIX

<b><u>GUARANTEED CONTENT</u></b>	<b>%W/W</b>
Water-Soluble Potassium Oxide (K <sub>2</sub> O)	25
Water-Soluble Magnesium Oxide (MgO)	10

**How to Apply the Medication**

<b><u>Plant Type</u></b>	<b><u>Application Period</u></b>	<b><u>In Leaf (with Gr/da 100 lt water)</u></b>	<b><u>In Drip Irrigation (with 1 ton water)</u></b>	<b><u>In Soil( Kg/da Per a Tree )</u></b>
Greenhouse Vegetables	Apply 4-5 times during the period starting from 4-5 leaf period of plants until harvest .		2-3 kg/decare	
Open Field Vegetables	Apply 4-5 times during the period starting from 4-5 leaf period of plants until harvest .		2-3 kg/decare	
Melon,Watermelon,Strawberry	Apply 4-5 times during the period starting from 4-5 leaf period of plants until harvest .		3-5 kg/decare	
Apple,Pear,Quince	Apply 3-4 times starting after fruit set with 20 days of intervals.	250-300 gr/100ltwater		
Peach,Berry,Cherry,Appricot	Apply 3-4 times starting after fruit set with 20 days of intervals.		2-4 kg/decare	
Grape,Pomegranate ,Fig	Apply 3-4 times starting after blooming with 20 days of intervals.			0,3 kg/decare
Banana	Apply 3-4 times starting after fruit set with 20 days of intervals.		3-5 kg/decare	0,5 kg/decare
Citrus,Olive,Tea	Apply 3-4 times starting after blooming with 20 days of intervals.		3-5 kg/decare	
Hazelnut,Walnut,Pistachio	During filling period	200-250 gr/100 lt water	2-3 kg/decare	0,5-1 kg/decare
Corn,Sunflower,Soy,Canola	From the period that height of plants reach 40-50 cm	200-250 gr/100 lt water	2-4 kg/decare	20-25 kg/decare
Cotton	From earing, until harvesting period	200-250 gr/100 lt water	2-3 kg/decare	20-25 kg/decare
Radish,Cabbage,Carrot,Celery, Cauliflower	Apply 1-2 times during the period starting from 4-5 leaf period of plants until harvest .	200-250 gr/100 lt water	2-3 kg/decare	20-25 kg/decare
Legumes, Forage Plants	Apply 1-2 times during the period starting from 4-5 leaf period of plants until harvest .	200-250 gr/100 lt water		20-25 kg/decare
Cereals(wheat,barley,oat, etc.)	From bolting period until latex beginning	200-250 gr/100 lt water		20-25 kg/decare
Sugar Beet,Potato,Paddy	Apply 1-2 times during the period starting from the formation of tuber, until the harvest.	250-300 gr/100 lt water	3-5 kg/decare	20-25 kg/decare
Onion,Garlic	Apply 1-2 times during the period starting from the formation of tuber, until the harvest.	200-250 gr/100 lt water	2-3 kg/decare	20-25 kg/decare
Ornamental Plants	Apply 2-3 times at 30 days intervals during growth periods	200-250 gr/100 lt water	2-3 kg/decare	20-25 kg/decare
Green Areas	Apply 2-3 times if needed	200-250 gr/100 lt water	2-3 kg/decare	20-25 kg/decare

License No:1484  
 Registration No:30236