

**EC FERTILIZER
BLENDED NPK FERTILIZER
PERFORMANS 18-18-18+ME**

<u>GUARANTEED CONTENT</u>	<u>% W/W</u>
Total Nitrogen (N):	18
Nitrate Nitrogen(NO3-N):	5
Ammonium Nitrogen(NH4-N)	7,7
Urea Nitrogen(NH2-N)	5,3
Water Soluble Phosphorus Pentoxide (P2O5)	18
Water Soluble Potassium Oxide(K2O)	18
EDTA Chelated Water Soluble Iron (Fe)	0,02
EDTA Chelated Water Soluble Manganese(Mn)	0,01
EDTA Chelated WATER Soluble Zinc (Zn)	0,002
Ph Range at which EDTA Chelating is stable Iron(Fe),Manganese(Mn),Zinc(Zn)	3-6

INTENDED USE, APPLICATION METHOD AND DOSAGE (Application In Soil)

<u>Plant Type</u>	<u>Application Period</u>	<u>Dosage</u>
Tomato, Pepper, Eggplant, Cucumber	1-2 applications per week after sowing until end of harvest	in leaves with 200-300 gr/100 lt water, 2-3 kg/da in roots
Melon-Watermelon	It must be applied 2-3 times during vegetation period.	in leaves with 300-350 gr/100 lt water, 3 kg/da in roots
Bean, Sweet pea	It must be applied 3-4 times at 10-day intervals, 2 weeks after sowing	200-300 gr/da in leaves, with 2-3 kg/da irrigation water
Cabbage-Cauliflower	It must be applied 3-4 times at 7-10 day intervals, 4-6 weeks after sowing	200-300 gr/da in leaves, with 2-3 kg/da irrigation water
Sugar beet, Carrot, Onion	It must be applied 2-3 times at 15-day intervals after second hoeing when number of leaves of the plants reaches 2-3	250-300gr/da in leaves, with 2-3 kg/da irrigation water
Strawberry	It must be applied 1-2 times before blooming, after blooming and fruit period.	in leaves with 200-250 gr/100 lt water, 2-3 kg/da in roots
Potato	It must be applied 3 times at stem elongation, budding and blooming phases.	in leaves with 200-300 gr/100 lt water, 3-4 kg/da in roots
Citrus fruits	It must be applied 3 times starting from before Blooming Period until the end of summer season.	3-4 kg/da in roots with 200-300 cc/100 lt water
Vineyard	It must be applied 3-4 times before blooming -blooming completed about 50% -seed formation starts	3-4 kg/da in roots with 200-250 cc/100 lt water
Peach, Apricot, Apple, Cherry Plum, Pear	It must be applied 2-3 times starting from spring season until the end of harvest period.	2-4 kg/da in roots with 200-250 cc/100 lt water
Wheat, Oat, Barley, Rye, Rough Rice	It must be applied twice at tillering season together with a Medication Against Weeds and earing period	in leaves with 200-300 gr/da 100 lt water
Cotton	At the phase of leave formation, blooming phase and at the phase of fruit formation.	3-4 kg/da in roots with 200-250 gr/da 100 lt water
Sunflower,Corn	When plants have 4-5 leaves.	3-4 kg/da in roots with 200-250 gr/da 100 lt water
Lentil, chickpea	It must be applied twice; before blooming and after blooming.	in leaves with 200-300 gr/da 100 lt water, 3-4 kg/da in roots
Pistachio	It must be applied 3-4 times 20-25 days after defloration and before nouaison .	in leaves with 200-300 gr/100 lt water, 2-3 kg/da in roots

Licence No:1484
Registration No:24321